

Care Gram

Medication Blood Levels

Some medications need blood levels checked. Testing the blood tells if there is too much or not enough of the medication in the body. After the results of the blood test, you may need to take more or less of the medication. Your doctor will let you know.

You must get your blood tested at just the right time. This will help the doctor make the best decision about your medications.

Find your medication listed below, and follow the directions to get your blood test done.

Drug Name	Drug Level Checked
<input type="checkbox"/> Amitriptyline (Tryptizol or Elavil)	12 hours after dose
<input type="checkbox"/> Digoxin (Lanoxin)	8-12 hours after dose
<input type="checkbox"/> Phenytoin (Dilantin or Epanutin)	Before next dose or 6 hours after dose
<input type="checkbox"/> Lithium	12 hours after dose
<input type="checkbox"/> Phenobarbital	Before next dose or 6 hours after dose
<input type="checkbox"/> Primidone (Mysoline)	Before next dose or 3 hours after oral dose
<input type="checkbox"/> Carbamazepine (Tegretol)	Before next dose or 6 hours after dose
<input type="checkbox"/> Theophylline	Before next dose or 6 hours after dose
<input type="checkbox"/> Valproic Acid (Depakote, Depakene or Epilim)	Before next dose

Thank you for choosing Licking Memorial Hospital Laboratory for your lab testing. We are open 24 hours a day, 365 days a year.

If you have any questions, please call (740) 348-4155.

This Care Gram is for information purposes only. It may become out-of-date over time. Ask a doctor or a healthcare professional about your specific condition.



**Licking Memorial
Health Systems**

1320 West Main Street • Newark, OH 43055
(740) 348-4000 • www.LMHealth.org

1616-0718
11/22/2010